



## Bedrock erosion landforms, Cantley, Québec

Leaders: David R. Sharpe, Geological Survey of Canada  
Guy Leduc, Douai, France

Mid-conference field trip: August 9, 2018



**Frontispiece:** aerial view of Cantley field site shows ~100 m scale linear rock drumlins and furrows: evidence of ice streaming? See more aerial views at this link: [https://youtu.be/wYp8noIc\\_Uk](https://youtu.be/wYp8noIc_Uk)

# CANQUA/AMQUA 2018

*Crossing borders in the Quaternary*

---

In memory of John Shaw

## **Abstract**

Several erosional forms on bedrock at Cantley, Quebec, differ from well-known glacial abrasion forms. The forms consist of obstacle marks, hollows, depressions, and channels, which are defined by sharp rims, smooth inner surfaces, divergent flow features, and remnant ridges. These forms are found on lee, lateral, and overhung rock surfaces. This assemblage of features is best explained by differential erosion produced by separation eddies along lines of flow reattachment. Rapid, sediment-laden, turbulent, subglacial melt-water flows likely produced the forms by corrasion and cavitation erosion.

Ice-abrasion forms, such as striations, and plucked forms such as gouges and crescentic fractures are also present at the Cantley site. Pitted forms, polishing, and carbonate precipitate are also present. The occurrence of abrasion, pitting, polishing, and lee-side carbonate precipitate with meltwater forms suggests that the meltwater flows were subglacial. Decoupling of abrading ice from its bed temporarily suspended glacial abrasion, whereas reattachment of ice to the bed may have led to the rounding of sharp edges and the production of striations superposed on the glaciofluvial forms.

The association of forms produced both by glaciofluvial erosion and ice abrasion suggests that the glacier was alternately lifted from, and reattached to, the bed during periodic subglacial floods. These floods may have affected the dynamics of the ice sheet, and depositional sequences related to high-energy meltwater outbursts were probably deposited in adjacent basins.


## Table of Contents

Abstract .....	2
Stop 1: Cantley bedrock erosional forms (s-forms) .....	4
Purpose .....	4
Description of site .....	4
Erosion forms .....	5
Striations and rat-tails .....	5
Obstacle marks and sichelwannen .....	6
Channels or Cavettos and Potholes .....	7
Other features .....	9
Cross-cutting forms .....	9
Interpretation .....	10
Glacial Abrasion .....	10
Corrasion in turbulent separated flow .....	10
Cavitation .....	11
Dissolution features .....	11
Discussion .....	11
Regional process implications .....	11
Key questions to consider during the Cantley field trip stop .....	12
References .....	12

## Stop 1: Cantley bedrock erosional forms (s-forms)

*David Sharpe, Geological Survey of Canada; Guy Leduc*

Location: Cantley Quebec, NTS 31G/12, UTM 18T 438721E 5048793N; 45° 35' N, 75° 47' W, ~ 190 m asl. Park in front of gate on the west side of Highway 307 (Fig. 1). Quarry owner is Gillies Tremblay; access at your own risk by writing to owner via the town at: 8, chemin River, Cantley, Quebec J8V 2Z9.



**Figure 1** Geology and digital terrain model of the Ottawa area showing fieldtrip stops and route. Inset provides setting of eastern Great Lakes. Related stops 2 and 3 are reported in Russell and Cummings, 2009.

### Purpose

The purpose of this stop is to draw attention to the significance and implications of erosion in glaciated terrain. Bedrock exposures within the quarry display subglacial erosional forms that are inferred to be mainly cut by powerful subglacial meltwater flow (Sharpe and Shaw, 1989) within a tributary valley of the Gatineau River (Fig. 1).

### Description of site

Bedrock within the quarry consists of Precambrian marble containing resistant granitic or volcanic clasts. The studied outcrop is streamlined parallel to valley orientation and to regional ~ north-south ice flow (Fig. 2). This site was deglaciated prior to 12 000 BP, but until recently, the described features were covered with gravel, sand, and mud deposited rapidly on subaqueous fans formed at the margin of the Champlain Sea (e.g. Rust, 1988; Sharpe, 1988), and below a marine limit of ~ 200 m asl.

Similar erosional forms occur on other outcrops in this poorly-exposed valley. Sculpted forms as observed on the bedrock surface diminish at higher elevation on adjacent outcrops. At the north end of the site, a 15 m thick sequence of glaciomarine sediment overlies the site and consist of a fining upwards sequence of sandy gravel and bedded sand overlain by silt and laminated clay. We provide an overview of the site features and interpretation with a drone-illustrated video with animation produced by Guy Leduc. View this link:

Cantley, Quebec - Monument of the Ice Age: [https://youtu.be/wYp8noIc\\_Uk](https://youtu.be/wYp8noIc_Uk)




Figure 2 Photograph of major outcrops at the Cantley quarry, Stop 1. R is rock drumlin, RR is remnant ridge. Outcrop forms are oriented parallel to the valley (~N-S) and in alignment with flow directions on small features in the quarry. Curved arrows indicate flow path that defines crescentic furrows. SF indicates sub-aqueous fan sediment that covered the rock surfaces. See person for scale.

## Erosion forms

Forms range in size from centimeter-scale striations and rat-tails, meter-scale obstacle marks, and forms the size of the outcrop, 50-100 m long, 5-10 m high (Fig. 2).

### *Striations and rat-tails*

Portions of the rock surface are planed-off and striated, such that the granitic and volcanic inclusions lie flush with the surface of the surrounding marble (Fig. 3). Small rat-tails are present in addition to striations. Rat-tails are positive forms and indicate differential erosion of the surrounding rock surface, rather than negative erosion of striations by a tool held in ice. Possible cavitation streaks, discontinuous grooves with pitted surfaces (Laguna-Camacho et al., 2013) were observed on some eroded rock surfaces (see Fig. 6b).






**Figure 3.** a) volcanic clast in Grenville marble is planed off on a striated rock surface displaying variable striae directions; flow bottomright to top left parallel to outcrop ~N-S orientation. b) rat-tail forms (r) (Prest, 1983), 1- 3 mm long, some with crescentic scours (cs) at the head and others without. Note pitted nature on the upflow (right side) of the dark rock inclusion. Flow right to left parallel to outcrop orientation. c) rat tails set within small shallow sculpted forms (s) oriented oblique to rat tails (r). Note: scale card is 8 cm long.

### ***Obstacle marks and sichelwannen***

The most common and distinctive erosion forms at the site are obstacle marks (Fig. 4), which have ridges in the lee of obstacles (Allen, 1982). These forms consist of a proximal, crescentic furrow wrapped around an upstanding obstacle, or resistant bedrock clast (Fig. 4b). The proximal furrow commonly has a sharp leading margin. The arms of the crescentic furrow extend downflow in a pair of furrows that become shallower and wider downflow (Fig. 4a, b).

The furrows are often smooth or less striated than adjacent surfaces outside the furrow (Fig 4a), and they can contain divergent flow features (Fig. 4b). Furrows also extend far downflow (Fig. 4c) to form longitudinal forms (Kor et al., 1991). Remnant ridges, which form behind the obstacle and between the furrows, may occur at several scales (compare Fig. 4a, c). An element (F; Fig. 4c), divergent from main flow, indicates secondary flow in the upflow furrow. Obstacle marks are common (Shaw, 1994) and appear to relate to complex flow.




**Figure 4** Obstacle marks (transverse s-forms): a) Sculpted forms diverge around dark inclusion (see 8 cms scale card at top) of obstacle within marble, set on a vertical rock face. Sculpted arms, or furrows (f), become wider and shallower downflow to the right, leaving a remnant ridge (r) in the lee of the obstacle. Note striation (s) or cavitation streak, and rat-tails on the rock surface. Note muschelbruche (mussel-shaped) scour (M). Flow left to right. b) Conceptual process model of obstacle mark formation under turbulent flow (Allen, 1971). c) A number of obstacle marks (O) occur on rock surface, some with large furrows (F) upflow of an obstacle and well-defined tapered remnant ridges (R). Note large, divergent rat-tail (rt) within the upflow furrow. Flow right to left parallel to outcrop orientation. Note Henri Richard for scale, lower left.

Other observed sculpted forms occur without obstacles; they are called *sichelwannen* (Ljugner, 1930) or *crescentic scours* (Dahl, 1965). These are now termed *s-forms* (Kor et al., 1991); they consist of transverse elements with furrows extending downflow around a central medial ridge (Fig. 2). At this scale, the remnant medial ridges can be considered to be rock *drumlins* (Fig. 2). A smaller, mussel-shaped transverse form (*muschelbruch*) occurs with sharp margins but without a medial ridge (see M, Fig. 4a).

### *Channels or Cavettos and Potholes*

Furrows that extend down flow (Fig. 4c) as longitudinal forms result in channels or *cavettos* (Dahl, 1965). They occur on vertical rock faces either as elongate troughs, or, as winding channels with tight

curves (Fig. 5). Cavettos are commonly meters long and centimeters to meters deep, and may crosscut other features (Fig. 5c). Most have vertical segments similar to truncated potholes (Gjessing, 1967). Vertical segments (Fig. 5b) are considered to be non-directional forms (Kor et al., 1991). At this site, rat-tails indicate that the eroding flow was only upward, unlike classical potholes, in which flow at the outside of the form is downward. Vertical flux implies flow under a very high hydrostatic pressure gradient.




**Figure 5** Furrows, cavettos and potholes: a) Poorly developed upflow furrow or crescentic scour (cs) wraps around obstacle (o) extending to lengthy longitudinal furrows (f, ~10 m) and remnant ridge (r). Flow right to left parallel to outcrop orientation. b) Cavetto or channel-like form with vertical orientation. Flow is upwards as indicated by rat-tails (arrow). This vertical structure had flow directed upwards rather than downward as in potholes. This indicates very high piezometric heads in the subglacial water system. Note John Shaw for scale. c) Complex pattern of longitudinal s-forms; furrows and cavettos. Flow left to right parallel to outcrop orientation. Note person lower right for scale.

### *Other features*

Interior surfaces of many s-forms are scored by light scratches (tool marks?). The scratches are short and are generally aligned with the direction of flow indicated by the divergent axes of troughs and associated rat-tails (Fig. 6). They also diverge, as the sculpted form becomes wider downflow.

In places, carbonate precipitate is visible on polished surfaces, particularly in the lee of obstacles and on remnant ridges (Fig. 6a). The precipitate likely formed in the late stages of flow (lee side occurrence) as sediments started to bury the landforms.

Pitting is visible on the upflow side of obstacles with remnant ridges (Fig. 6b). This appears to be incipient development of upflow furrows, perhaps as an emerging rock ridge is exposed to accelerated erosion.



**Figure 6** Other erosional features: a) Short, light scratches occur in alignment with the axes of furrows (f) and associated remnant ridge (r) surfaces. Carbonate precipitate (C) is also found on polished surfaces in places, particularly in the lee of obstacles (O) and on remnant ridges (see 8 cm card). b) Pitting (P) occurs on the upflow side of an obstacle (o) with a remnant ridge (RR) (see also Fig. 5b). Note striations (s) adorn this surface and form at slight cross-cutting angles; note possible discontinuous cavitation streak (c). c) Striations (s), set at an oblique angle, occur on top of transverse s-form, hairpin scour (HS). Striae oblique to hairpin scour record the glacier sole re-attaching to the sculpted rock surface with little erosional force.

### *Cross-cutting forms*

Numerous cross-cut relationships involving striations and sculpted forms are visible at the Cantley site (Fig. 4c, 5b, c, 6c). For example, striations occur as ornamentations, set at an oblique angle, on top of

a transverse s-form (sichelwannen). In some forms, rat-tails readily show divergent flow within s-forms (Fig. 5c).

## **Interpretation**

### ***Glacial Abrasion***

From the evidence of striations on the surfaces of some sculpted marks, it is concluded that the erosional forms at Cantley were created subglacially. Striations and planed-off inclusions indicate glacial abrasion. Such abrasive planing is attributed to erosion by debris in the bed of flowing glacial ice (Hallet, 1981). There are fewer planed erosion surfaces than surfaces carrying sculpted forms (s-forms). S-forms, in contrast to striations, likely relate to differential erosion expected in flow systems with secondary, turbulent structures (Allen, 1971). Alternatively, the possibility of glacial formation of obstacle marks considers laminar streaming of debris-rich glacial ice around an obstacle (Boulton, 1974). However, because ice is incompressible, there is no increase in debris concentration (Boulton, 1979) and no enhanced erosion. Gjessing (1965) proposed that, although some s-forms result from fluvial erosion and cavitation, others are a product of erosion by a subglacial slurry of saturated till. Others suggest that the saturated till, as a deforming bed, was responsible for erosion of streamlined rock forms (e.g. O' Cofaigh et al., 2005). The laminar, creeping motion of slurry or deforming bed flow does not explain hairpin scours (Fig. 6 c) and related transverse erosional marks with a dearth of striations.

### ***Corrasion in turbulent separated flow***

Cantley obstacle marks, and other transverse s-forms, are thought to result from glaciofluvial processes (Shaw, 1988; Sharpe and Shaw, 1989; Kor et al., 1991; Munro-Stasiuk et al., 2005). S-forms associated with obstacles are likely formed by flow separation and horseshoe vortices (Fig. 4c), resulting from a vertical pressure gradient generated at the up-stream face of the obstacle in turbulent flow (Shaw, 1988; 1994). This gradient sets up secondary flow and a pair of oppositely rotating vortices (Fig. 4c). These secondary flows reattached to the bed where high velocity fluid approached at a high angle and caused maximum erosion (Allen, 1971, 1982). The crescentic furrow cut around the leading side of the obstacle and the paired furrows extending downflow are likely products of a horseshoe vortex (Dargahi, 1990; Shaw, 1994). These vortices expanded rapidly and thus, became reduced in intensity and erosional power downstream. As a consequence, the furrows become broader and shallower downflow (Fig. 4b, c), and, rat tails, the remnant (rock) ridges between the furrows, becomes narrower and lower. Sichelwannen similarly relate to flow separation and horseshoe vortices (Kor et al., 1991), whereas muschelbruch (Fig. 4a, b) likely relate to impingement of low-angle vortices on the bed (Shaw, 1988). Thus, obstacle marks clearly relate to the generation of horseshoe vortices (Fig. 4b) in turbulent flow, a process that is common in both nature (Doumani, 1967; Shaw, 1994) and in engineering fluid dynamic experiments (Paik et al., 2010).

Some s-forms that are oblique to the general flow direction indicated by a longitudinal element are preserved due to their protected location within a furrow. Most appear to represent secondary flows related to vortices within a primary flow. This type of mechanical erosion at Cantley is considered to be glaciofluvial corrasion and can occur by direct fluid stressing, or, it can be related to erosion by tools carried in flowing water (Allen, 1982).

Smaller scale forms are identical to some sculpted fluvial forms in terrain subject to flooding in Australia (Baker and Pickup, 1986). Some larger forms are similar to forms identified within the Channeled Scablands of Washington State (Bretz, 1925; Baker, 1978). These comparisons support the interpretation of formation by water erosion. Allen (1971, 1982) also verified by way of flume experiments that erosional forms such as these observed at Cantley can be produced by high-velocity, separated, turbulent fluid flow.

### ***Cavitation***

Cavitation is the rapid formation and collapse of bubbles due to local pressure gradients in turbulent flow. It becomes a viable erosion process where turbulent flow has high enough velocity (~5-10 m/s; Hjulstrom, 1935; Richardson and Carling, 2005). Damaging shock waves and violent jets of water score turbine blades, and can loosen grains in rocks, pit massive rocks (Dular et al., 2004), while creating longitudinal streaks such as may be inferred in places at Cantley (Fig. 6c).

### ***Dissolution features***

There is some evidence of dissolution of Precambrian marble at Cantley. There are, however, few dissolution features, and carbonate precipitation indicates a non-dissolution regime following main subglacial erosion. It does not appear to be a dominant process, despite the suggestion of Hillaire-Marcel (2005) that the site 'represents the mixed influence of mechanical abrasion by basal ice and of dissolution features due to high pressure subglacial water channeling'. Dissolution features probably relate to minor post-glacial modification of common sculpted forms that has taken place on marble surfaces following removal of sand and gravel from the quarry rock surface in recent years.

### **Discussion**

The outcrops at Cantley display s-forms (e.g. obstacle marks, furrows, and cavettos) with sharp rims, divergent flow features, and remnant ridges. They also show ice abrasion forms, striations, and plucked forms such as crescentic fractures. The occurrence of abrasion, pitting, polishing, and carbonate precipitate with meltwater forms suggests that the meltwater flows were subglacial. Lifting of ice from its bed by fast-flowing meltwater suspended glacial abrasion. When ice settled back on the bed, as the meltwater flow subsided, abrasion resumed, rounding sharp edges and lightly striating rock faces, in places at oblique directions.

The association of forms produced both by glaciofluvial erosion and ice abrasion suggests that the glacier was lifted from and let down on the bed during subglacial floods. The assemblage of sculpted features at Cantley is best explained by differential erosion produced by strong vortices. Rapid, sediment-laden, turbulent, subglacial meltwater flows likely produced most forms by corrasion and cavitation erosion (Sharpe and Shaw, 1989). Depositional sequences (sand and gravel) related to these high-energy meltwater outbursts were probably deposited on subaqueous fans in the adjacent Champlain Sea basin (e.g. Rust, 1988; Sharpe, 1988; Cummings et al., 2011).

### **Regional process implications**

Water-sculpted erosion forms imply subglacial bed conditions with little frictional resistance in the areas of subglacial meltwater flow at the time of such discharge (Shoemaker, 1992). The large inferred discharge rates also require meltwater storage, likely in subglacial reservoirs (e.g. Alley et al., 2006; Evatt et al., 2006) and supraglacial lakes (Zwally, 2002), up-ice from the Cantley site. Such stored

subglacial water may have occurred in Hudson Bay or in any bed-roughness element on the scale of 5 km or more (e.g. modern day lake basin) that are found in abundance on the Canadian Shield (Shoemaker, 1992). Separation of the glacier from its bed by a subglacial meltwater sheet, as in some Icelandic floods, also involves minimal basal resistance to ice flow (Shoemaker, 1992; 1999), and a flat ice sheet is expected to cover such discharge and storage areas (Wingham et al., 2006). The importance given here to meltwater events correspond to that applied by others to subglacially deforming till beds (e.g. Boulton and Hindmarsh, 1987; O’Cofaigh et al., 2005). Both process models imply low basal shear stresses, a relatively flat ice-sheet profile, and minimum ice volumes for a given ice-sheet radius. It is important to note that, in the meltwater explanation, these conditions may occur even where the substrate is bedrock. The rapid discharge of meltwater interpreted here is expected to have been accompanied by accelerated ice flow, perhaps surging (Kamb et al, 1985), or ice streaming (Bell et al., 2007). If ice-sheet profiles are to be credibly reconstructed, it becomes critical that subglacial meltwater forms be mapped and their timing be assessed.

### **Questions to consider during the Cantley field trip stop**

1. Are the elongate rock ridges in the aerial view of Cantley, mega-scale lineations?
2. How do the aerial view rock features compare with those viewed on the outcrop?
3. Are sculpted erosion forms (s-forms) distinguishable from ice flow indicators (striations etc.)?
4. Should s-forms be mapped and recorded on maps similar to the way that ice flow indicators (striations) are shown?
5. Is it possible that s-forms and ice flow forms record different elements of the same event, or closely timed event?
6. If s-forms are present across a region, is it reasonable to interpret a regional event(s) as is the case of with striation mapping.
8. Are ice streams required to explain any flow features at Cantley?
9. What features at Cantley might indicate rapid flow?
10. Is there sedimentological evidence for the release of large volumes of stored meltwater that scoured the Cantley site?

### **References**

- Allen, J.R.L., 1971. Transverse erosional marks of mud and rock: Their physical basis and geological significance: *Sedimentary Geology*, 5: 167-185.
- Allen, J.R.L., 1982. *Sedimentary structures: Their character and physical basis*. Volume 11; New York, Elsevier Scientific Publishing Co., p. 253-291.
- Alley, R.B., Dupont, T.K., Parizek, K.B.R., Anandakrishnan, S., Lawson, D.E., Larson, G.L., and Evenson, E.B., 2006. Outburst flooding and the initiation of ice-stream surges in response to climate cooling: a hypothesis. *Geomorphology*, 75: 76-89.
- Baker, V. R., 1978. The Spokane Flood controversy and the Martian outflow channels. *Science*, 202: 1249-1256.
- Baker, V. R., and Pickup, G., 1987. Flood geomorphology of the Katherine Gorge, Northern Territory. *Australia: Geological Society of America Bulletin*, 98: 635-646.
- Bell, R.E., Studinger, M., Shuman, C., Fahnstock M. A. and Joughin I., 2007. Large subglacial lakes in East Antarctica at the onset of fast-flowing ice streams. *Nature*, 445: 904-907.

- Boulton, G.S., 1974. Processes and patterns of glacial erosion, in Coates, D.R., ed. *Glacial Geomorphology*. Binghamton, New York, State University of New York, p. 41-87.
- Boulton, G.S., 1979. Processes of glacial erosion on different substrata. *Journal of Glaciology* 23, 15-38. (with comment by Barclay Kamb).
- Boulton, G. S., and Hindmarsh, R.C.A., 1987. Sediment deformation beneath glaciers: Rheology and geological consequences: *Journal of Geophysical Research*, 92(B9): 9059-9082.
- Bretz, J.H., 1925. The Spokane flood beyond the Channeled Scabland. *Journal of Geology* 33, 97-115.
- Cummings, D.I., Pugin, A., Pullan, S., Gorrell, G., Ponomarenko, D., Hunter, J., Russell, H.A.J., and Sharpe, D.R., 2011. Geology of muddy glaciated-basin fill, with a focus on esker sedimentation. *Geological Society of America Bulletin*.  
<http://gsabulletin.gsapubs.org/content/early/2011/01/26/B30273.1.abstract>
- Dahl, R., 1965. Plastically sculptured detail forms on rock surfaces in northern Nordland, Norway. *Geografiska Annaler*, 74A: 83-140.
- Dargahi, B., 1990. Controlling mechanism of local scouring. *Journal of Hydraulic Engineering* 116, 1197-1214.
- Doumani, G.A., 1967. Surface structures in snow. In. Oura H. (Ed.) *Physics of Snow and Ice*. Institute of Low Temperature Science. Hokkaido University of Japan, 1119-1136.
- Dular, M., Bachert, B., Stoff, B., Sirok, B., 2004. Relationship between cavitation structures and cavitation damage. *Wear* 257, 1176-1184.
- Evatt, G.W., Fowler, A.C., Clark, C.D. and N. R. J. Hulton. 2006. Subglacial floods beneath ice sheets. *Philosophical Transactions of the Royal Society*, 364, 1769-1794.
- Gjessing, J., 1965. On "plastic scouring" and subglacial erosion. *Norsk Geografisk Tidsskrift* 20, 1-37.
- Gjessing, J. 1967. Potholes in connection with plastic scouring forms: *Geografiska Annaler*, 48A(2): 178-187.
- Hallet, B. 1981, Glacial abrasion and sliding: Their dependence on the debris concentration in basal ice: *Annals of Glaciology*, 2: 23-28.
- Hillarie-Marcel, C., 2005. Cover photo and caption, *Canadian Journal of Earth Sciences*, 42(12).
- Hjulström, F., 1935. Morphological activity of rivers as illustrated by the river Fyris: *Bulletin of the Geological Institute of Uppsala*, 25: 221-527.
- Kamb, B., Raymond, C.F., Harrison, W.D., Engelhardt, H., Echelmeyer, K.A., Humphrey, N., Arugman, M.M., and Pfetter, T., 1985. Glacial surge mechanism: 1882-1983 surge of Variegated Glacier, Alaska: *Science*, 227: 469-479.
- Kor, P., Shaw, J., and Sharpe, D.R., 1991. Erosion of bedrock by subglacial meltwater, Georgian Bay, Ontario: a regional view. *Canadian Journal of Earth Sciences*, 28(4): 623-642.
- Laguna-Camacho, J.R., Lewis, R., Vite-Torres, M., Méndez-Méndez, J.V., 2013. A study of cavitation erosion on engineering materials. *Wear* 301, 467-476.
- Ljungner, E., 1930. Spaltetentektonik und Morphologie der schwedischen skager-küste. III:I. *Bulletin of the Geological Institutes of the University of Uppsala* 21, 255-475.
- Munro-Stasiuk, M. J., Fisher, T. G., and Nitzsche, C. R., 2005. The origin of the western Lake Erie grooves, Ohio; implications for reconstructing the subglacial hydrology of the Great Lakes sector of the Laurentide ice sheet. *Quaternary Science Reviews*, 24: 2392-2409.
- O'Cofaigh, C., Dowdeswell, J.A., Allen, C.S., Hiemstra, J.F., Pudsey, C.J., Evans, J. and Evans, D.J.A., 2005. Flow dynamics and till genesis associated with a marine-based Antarctic palaeo-ice stream. *Quaternary Science Reviews*, 24: 709-740.
- Paik, J., Escauriaza, C., Sotiropoulos, F., 2010. Coherent structure dynamics in turbulent flows past in-stream structures: some insights gained via numerical simulation. *Journal of Hydraulic Engineering* 136 doi: 10.61/(ASCE)HY.1943-7900.000089

- Prest, V., .K. 1983. Canada's heritage of glacial features. Geological Survey of Canada, Miscellaneous Report 28.
- Richardson, K. and P.A., Carling, 2005. A typology of sculpted forms in open bedrock channels. Geological Society of America, Special Paper 393, 108p.
- Russell, H.A.J., and D. Cummings, 2009. Deglaciation of the Champlain Sea basin, eastern Ontario. 72nd FRIENDS OF THE PLEISTOCENE reunion, June 6–7, 2009, Ottawa, Ontario.
- Rust, B.R. 1988. Ice-proximal deposits of the Champlain Sea at south Gloucester, near Ottawa, Canada. In: The Late Quaternary Development of the Champlain Sea Basin (Ed N.R. Gadd), 35, p. 37-45. Geological Association of Canada, Special Paper, St John's, NF.
- Sharpe, D.R. 1988. Glaciomarine fan deposition in the Champlain Sea. In: The Late Quaternary Development of the Champlain Sea Basin (Ed N.R. Gadd), 35, p. 63-82. Geological Association of Canada, Special Paper, St. John's, NF.
- Sharpe, D.R and Shaw, J. 1989. Erosion of bedrock by subglacial meltwater, Cantley, Sharpe, D.R., 2005. Comments on “Paleohydrology of the last outburst flood from glacial Lake Agassiz and the 8200 B.P. cold event”: Quaternary Science Reviews, 28: 1529-1532.
- Shaw, J., 1988. Subglacial erosion marks, Wilton Creek, Ontario. Canadian Journal of Earth Sciences, 25: 1256-1267.
- Shaw, J., 1994. Hairpin erosional marks, horseshoe vortices and subglacial erosion: Sedimentary Geology, 91: 269-283.
- Shoemaker, E.M., 1992. Water sheet outburst floods from the Laurentide ice sheet; Canadian Journal of Earth Sciences, 29: 1250-1264.
- Shoemaker, E.M., 1999. Subglacial water-sheet floods, drumlins and ice-sheet lobes. Journal of Glaciology, 4: 323-329.
- Wingham, D.J., Siegert, M.J., Shepherd, A.P. and Muir, A.S., 2006, Rapid discharge connects Antarctic subglacial lakes. Nature, v. 440, p. 1033–1036.
- Zwally, H.J., Abdlati, W., Herring, T., Larson, K., Saba, J., Steffenn, K., 2002. Surface melt induced acceleration of Greenland ice sheet flow. Science 297, 218-222